

*Microbial Science Archives (ISSN: 2583-1666)*

An International Research Journal

Journal homepage: [www.microbial.sciencearchives.org](http://www.microbial.sciencearchives.org)

<https://doi.org/10.47587/MSA.2026.6202>

[Review Article](#)



## Plant-Microbe Interactions: A Review of Current Research and Future Directions in Sustainable Agriculture

Odoh Emeka Hyginus<sup>1</sup> , Eze Constance Nonye<sup>2</sup> , Jude, Egeonu Vitus<sup>3</sup> , Hamisu Shuaibu Idris<sup>4</sup> , Moses Adondua Abah<sup>5,6</sup>  , Micheal Oladosu Abimbola<sup>6,7</sup>  and Ochuele Dominic Agida<sup>5,6</sup> 

<sup>1</sup>*Department of Biology, Faculty of Arts and Sciences, Georgia State University, USA*

<sup>2</sup>*Department of Biological Sciences, Southern Illinois University, Edwardsville, Illinois United States.*

<sup>3</sup>*Department of Microbiology/Biotechnology, Faculty of Sciences, Federal University Dutse, Jigawa State, Nigeria*

<sup>4</sup>*Department of Microbiology/Biotechnology, Faculty of Life Sciences, Federal University Dutse, Dutse, Jigawa State, Nigeria*

<sup>5</sup>*Department of Biochemistry, Faculty of Biosciences, Federal University Wukari, Taraba State, Nigeria*

<sup>6</sup>*ResearchHub Nexus Institute, Nigeria*

<sup>7</sup>*Department of Chemical Sciences, Faculty of Science, Anchor University, Ayobo, Lagos State, Nigeria*

(✉) **Corresponding Author:** Moses Adondua abah; [m.abah@fuwukari.edu.ng](mailto:m.abah@fuwukari.edu.ng)

Received: April 28, 2026/ Revised: May 29, 2026/Accepted: June 2, 2026

### Abstract

Plant–microbe interactions play a fundamental role in maintaining plant health, soil fertility, and agricultural productivity. In recent years, growing environmental concerns associated with excessive use of chemical fertilizers and pesticides have intensified interest in sustainable agricultural approaches driven by beneficial microorganisms. This review examines the current understanding of plant–microbe interactions and their emerging significance in sustainable agriculture. It discusses the diversity of plant-associated microorganisms, including nitrogen-fixing bacteria, mycorrhizal fungi, plant growth-promoting rhizobacteria, endophytes, and biocontrol agents, highlighting their roles in nutrient cycling, plant growth enhancement, stress tolerance, and disease suppression. The review further explores the molecular and biochemical mechanisms underlying these interactions, with emphasis on signaling pathways, phytohormone production, and microbial-mediated resistance responses. Recent advances in genomics, metagenomics, microbiome engineering, synthetic biology, and artificial intelligence are also examined as transformative tools for understanding and optimizing plant–microbe systems. In addition, the review evaluates practical applications of microbial technologies in biofertilizers, biostimulants, and sustainable crop management strategies aimed at improving soil health and agricultural resilience under changing climatic conditions. Despite significant progress, challenges such as environmental variability, inconsistent field performance, formulation limitations, and regulatory constraints continue to hinder large-scale adoption. The review concludes by highlighting future research priorities focused on climate-smart agriculture, next-generation microbial consortia, and precision microbiome management. Overall, plant–microbe interactions represent a promising foundation for developing environmentally sustainable, resilient, and productive agricultural systems capable of supporting global food security.

**Keywords:** Plant–microbe interactions, Sustainable agriculture, Rhizosphere microbiome, Biofertilizers, Rhizobacteria and Microbiome

## Introduction

Modern agriculture faces increasing pressure to meet the growing global demand for food while addressing environmental degradation, declining soil fertility, biodiversity loss, and climate change. Although the intensive use of chemical fertilizers and pesticides has significantly improved crop productivity, these practices have also generated serious ecological concerns, including soil contamination, nutrient imbalance, greenhouse gas emissions, and disruption of natural microbial communities. Consequently, sustainable agricultural approaches that enhance productivity while preserving environmental quality are receiving global attention. Among the most promising biological strategies is the utilization of plant–microbe interactions, which play essential roles in maintaining soil health, supporting plant growth, and improving agricultural sustainability (Nadarajah and Abdul Rahman, 2023). Increasing evidence indicates that plant-associated microorganisms are fundamental to ecosystem functioning and may provide environmentally friendly alternatives to synthetic agricultural inputs (Compant *et al.*, 2025).

Plant–microbe interactions involve diverse associations between plants and microorganisms occurring within the rhizosphere, phyllosphere, and plant endosphere. These interactions may be beneficial, neutral, or pathogenic depending on the microbial species involved and environmental conditions. Beneficial microorganisms such as nitrogen-fixing bacteria, arbuscular mycorrhizal fungi, endophytes, and plant growth-promoting rhizobacteria (PGPR) contribute significantly to plant growth and stress adaptation (Liu-Xu *et al.*, 2024). In the rhizosphere, microorganisms interact closely with plant roots and participate in nutrient cycling, organic matter decomposition, and soil aggregation. Many beneficial microbes improve nutrient availability by solubilizing phosphorus, fixing atmospheric nitrogen, and mobilizing essential micronutrients. In addition, microbial communities produce phytohormones, enzymes, and bioactive metabolites that stimulate root development, improve water uptake, and enhance tolerance to abiotic stresses such as drought, salinity, and extreme temperatures (Tharanath *et al.*, 2024). These microbial associations also contribute to disease suppression through antibiosis, competition, and induced systemic resistance, thereby reducing dependence on chemical pesticides and supporting sustainable crop protection (Alzate Zuluaga *et al.*, 2024).

Scientific understanding of plant–microbe interactions has evolved substantially from early observations of visible symbiotic associations to advanced molecular investigations of complex microbial communities. Earlier studies mainly focused on biological nitrogen fixation and mycorrhizal symbioses because of their direct effects on plant nutrition and growth. However, recent advances in genomics, metagenomics, transcriptomics, metabolomics, and bioinformatics have transformed understanding of plant

microbiomes and their ecological functions in agricultural systems (Zhu *et al.*, 2024). These technologies enable the identification of previously unculturable microorganisms and provide deeper insights into microbial communication, signaling pathways, and plant immune responses. Furthermore, microbiome engineering and synthetic biology are creating new opportunities for designing microbial consortia with improved agricultural functions. Artificial intelligence and precision agriculture technologies are also being integrated into microbiome research to optimize microbial applications and predict plant–microbe responses under varying environmental conditions (Compant *et al.*, 2025).

The increasing demand for environmentally sustainable agricultural systems has accelerated the development of microbial-based products such as biofertilizers, biostimulants, and biocontrol agents. These products are increasingly recognized for their ability to improve nutrient use efficiency, enhance soil fertility, and support environmentally friendly crop production practices (Alzate Zuluaga *et al.*, 2024). Beneficial microorganisms also contribute to ecosystem restoration, carbon sequestration, and soil structural stability, making them important components of regenerative and climate-smart agriculture. Endophytic microorganisms, in particular, have attracted significant attention because of their capacity to improve plant adaptation to changing environmental conditions and enhance crop productivity under climate stress (Sena *et al.*, 2024). Despite these advantages, practical implementation of microbial technologies still faces limitations, including inconsistent field performance, environmental variability, poor microbial survival, formulation difficulties, and regulatory challenges (Tharanath *et al.*, 2024).

Given the growing need for sustainable and resilient agricultural systems, understanding the diversity, mechanisms, and applications of plant–microbe interactions has become increasingly important in scientific research and agricultural practice. Beneficial microbial communities provide significant opportunities for improving crop productivity while reducing the environmental impacts associated with intensive farming practices. Therefore, this review provides a comprehensive overview of current research on plant–microbe interactions and their significance in sustainable agriculture. The review examines the major categories of plant-associated microorganisms, the mechanisms underlying microbial-mediated plant growth and protection, and the practical applications of microbial technologies in modern agriculture. In addition, recent technological innovations, emerging research trends, major limitations, and future perspectives associated with microbial-assisted agriculture are discussed to highlight their potential contributions toward achieving sustainable global food security.

## Historical Background and Research Progress

### Early Discoveries

The scientific understanding of plant–microbe interactions has evolved through centuries of agricultural observation and experimental research. Early farming communities recognized the importance of fertile soils and organic residues in improving crop productivity, although the microbial mechanisms responsible for these effects remained unknown. The formal study of plant-associated microorganisms began during the nineteenth century with the emergence of microbiology and plant physiology as scientific disciplines. One of the earliest major breakthroughs was the discovery of biological nitrogen fixation in legumes, where symbiotic bacteria later identified as *Rhizobium* species colonize root nodules and convert atmospheric nitrogen into plant-available forms (Oldroyd and Dixon, 2014). This discovery provided one of the first clear demonstrations of mutualistic relationships between plants and microorganisms and significantly transformed agricultural and soil fertility research. Subsequent investigations during the late nineteenth and early twentieth centuries revealed the ecological importance of mycorrhizal fungi in plant nutrition and soil functioning. Researchers demonstrated that arbuscular mycorrhizal fungi establish intimate associations with plant roots, enhancing phosphorus uptake, water absorption, and plant adaptation to nutrient-deficient soils (Genre *et al.*, 2020). During the same period, studies on microbial plant diseases contributed to the establishment of plant pathology as an independent scientific field. Scientists recognized that microorganisms could exert both beneficial and harmful effects on plants, thereby expanding scientific understanding of host–microbe interactions in agricultural ecosystems (Compant *et al.*, 2021).


Another important milestone was the introduction of the rhizosphere concept by Lorenz Hiltner in 1904. The rhizosphere was described as the biologically active soil region surrounding plant roots, where intense interactions occur between plants, microorganisms, and soil components (Philippot *et al.*, 2013). This concept became central to understanding microbial diversity, nutrient cycling, root exudation, and soil biochemical processes. Further studies revealed that root-associated microorganisms contribute significantly to phosphorus solubilization, nitrogen fixation, phytohormone production, and pathogen suppression (Backer *et al.*, 2018). These discoveries laid the foundation for the development of microbial inoculants, biofertilizers, and biological control strategies currently applied in sustainable agriculture. By the mid-twentieth century, advances in soil microbiology and microbial ecology provided deeper insights into microbial-mediated nutrient cycling and plant health. Researchers demonstrated that microbial communities influence soil fertility, organic matter decomposition, carbon sequestration, and disease resistance within agricultural systems (Finkel *et al.*, 2017). The discovery of endophytic

microorganisms inhabiting internal plant tissues without causing disease further expanded understanding of plant-associated microbial diversity. Endophytes were subsequently found to promote plant growth, improve stress tolerance, and enhance resistance against pathogens and environmental stressors. These scientific developments established the foundation for integrating microbial ecology into modern sustainable agricultural systems.

### Advances in Molecular and Microbiome Research

The emergence of molecular biology and high-throughput sequencing technologies revolutionized plant–microbe interaction research during the late twentieth and early twenty-first centuries. Traditional culture-dependent methods were limited because a large proportion of plant-associated microorganisms could not be cultivated under laboratory conditions. However, the development of molecular tools such as polymerase chain reaction (PCR), DNA fingerprinting, and next-generation sequencing enabled researchers to analyze microbial communities directly from environmental samples without cultivation (Berg *et al.*, 2020). These technologies significantly expanded understanding of microbial diversity and revealed that plants host highly complex and dynamic microbiomes that strongly influence plant physiology, immunity, and ecosystem functioning. Modern omics technologies, including metagenomics, transcriptomics, proteomics, and metabolomics, have further advanced understanding of microbial community structure and function in agricultural ecosystems. These approaches allow detailed investigation of microbial genes, metabolic pathways, signaling networks, and microbial interactions within the plant holobiont (Trivedi *et al.*, 2020). Research has demonstrated that beneficial microorganisms contribute not only to nutrient acquisition and plant growth but also to resistance against drought, salinity, heavy metals, and pathogen attack (Müller *et al.*, 2016). In addition, environmental factors such as soil type, climate, agricultural practices, and host genotype have been identified as key determinants influencing microbiome composition and function (Zhu *et al.*, 2024).

Recent advances in synthetic biology, microbiome engineering, and computational biology are rapidly transforming plant–microbe interaction research from descriptive ecological studies into predictive and application-oriented sciences. Artificial intelligence, machine learning, and bioinformatics tools are increasingly being used to analyze complex microbiome datasets, predict microbial behavior, and optimize microbial inoculant performance in agricultural systems (Levy *et al.*, 2018). Furthermore, synthetic microbial communities are being designed to improve nutrient use efficiency, disease resistance, and crop resilience under environmental stress conditions (Vorholt *et al.*, 2017). These technological innovations are creating new opportunities for precision microbiome management and sustainable agricultural practices capable of supporting future global food security.



**Figure 1. Historical evolution of plant–microbe interactions from early microbial life to modern plant microbiome research.** This figure illustrates the evolutionary progression of plant–microbe interactions from the emergence of early microbial life to the development of complex plant microbiomes in modern agricultural systems. It highlights major milestones in microbial evolution, plant colonization of land, symbiotic associations, and advances in microbiome research that have contributed to current understanding of sustainable plant–microbe relationships.

Source: [Shelake et al. \(2026\)](#)

### Types of Plant–Microbe Interactions

Plant–microbe interactions are complex biological relationships established between plants and diverse groups of microorganisms inhabiting different plant-associated environments. These interactions occur mainly within the rhizosphere, phyllosphere, and endosphere, where microorganisms influence plant growth, nutrient acquisition, stress tolerance, immunity, and overall ecosystem functioning. Depending on the ecological outcomes for the host plant, plant–microbe interactions are generally classified as beneficial, neutral, or pathogenic. The balance among these interaction types largely determines plant health, soil fertility, microbial diversity, and agricultural productivity ([Berendsen et al., 2012](#)). Understanding these interactions is therefore essential for developing sustainable agricultural systems that maximize beneficial microbial activities while minimizing plant diseases and environmental degradation.

### Beneficial Interactions

Beneficial plant–microbe interactions involve microorganisms that positively influence plant growth, nutrient acquisition, and environmental adaptation. Major beneficial microbes include plant growth-promoting rhizobacteria (PGPR), arbuscular

mycorrhizal fungi, nitrogen-fixing bacteria, and fungal endophytes. These microorganisms enhance plant productivity through mechanisms such as nutrient mobilization, phytohormone production, biological nitrogen fixation, and pathogen suppression ([Santoyo et al., 2021](#)). Arbuscular mycorrhizal fungi improve phosphorus uptake, water absorption, and nutrient acquisition, especially under nutrient-deficient conditions ([Begum et al., 2019](#)), while nitrogen-fixing bacteria enrich soil fertility by converting atmospheric nitrogen into plant-available forms. Beneficial microorganisms also strengthen plant tolerance to abiotic stresses including drought, salinity, temperature fluctuations, and heavy metal toxicity. Many rhizosphere microbes stimulate root growth and regulate plant hormones, while others produce antimicrobial compounds that enhance plant defense and induce systemic resistance against pathogens ([Bais et al., 2005](#)). In addition, microbial consortia often exhibit synergistic effects that improve nutrient use efficiency and crop productivity in sustainable agricultural systems ([Santoyo et al., 2021](#)). Fungal endophytes further contribute to plant growth, stress adaptation, and disease resistance through the production of bioactive metabolites, making them valuable for biofertilizers and eco-friendly crop protection strategies ([Lugtenberg et al., 2016](#)).

### Neutral Interactions

Neutral plant–microbe interactions occur when microorganisms colonize plant-associated environments without exerting obvious positive or negative effects on plant growth or health. These microorganisms often exist as part of the natural plant microbiome and contribute to microbial diversity and ecological stability within plant ecosystems. Although neutral microbes may not directly influence plant productivity under normal environmental conditions, they can become functionally important during environmental disturbances or shifts in microbial community structure (Compant *et al.*, 2022). Some neutral microorganisms contribute indirectly to nutrient cycling, microbial competition, and maintenance of microbiome balance within the rhizosphere and plant tissues. Recent studies indicate that neutral microbial populations contribute significantly to functional redundancy within microbial communities, thereby improving ecosystem resilience under fluctuating environmental conditions (Niu *et al.*, 2017). Functional redundancy allows different microbial species to perform similar ecological functions, ensuring the stability of plant-associated microbial communities during environmental stress. Furthermore, rhizosphere microbial diversity is strongly influenced by soil properties, salinity, cropping systems, and root exudates, all of which affect microbial composition and ecological interactions within agricultural systems (Chandra *et al.*, 2020; Wang *et al.*, 2022). Consequently, neutral interactions are increasingly recognized as important components of plant microbiome ecology and sustainable agriculture.

### Pathogenic Interactions

Pathogenic plant–microbe interactions involve microorganisms that invade plant tissues and disrupt normal physiological and metabolic functions, resulting in disease development and reduced agricultural productivity. Plant pathogens include bacteria, fungi, viruses, oomycetes, and nematodes capable of causing severe crop losses worldwide. These pathogenic microorganisms infect plants through diverse mechanisms involving tissue colonization, secretion of toxins, degradation enzymes, and suppression of host immune responses (Xin *et al.*, 2018). Diseases caused by microbial pathogens significantly threaten global food security and agricultural sustainability. Plant pathogenic microorganisms

interact with host plants through highly sophisticated molecular signaling pathways. Successful pathogens often secrete effector proteins that interfere with plant immune systems and facilitate microbial colonization within plant tissues. In response, plants activate defense mechanisms including pattern-triggered immunity, effector-triggered immunity, oxidative burst responses, and antimicrobial compound production (Ngou *et al.*, 2021). Environmental conditions such as temperature, moisture, nutrient availability, and soil microbial composition strongly influence pathogen virulence and disease severity. Understanding pathogenic interactions is therefore essential for improving disease management strategies, developing resistant crop varieties, and reducing the economic impact of plant diseases in agriculture.

### Rhizosphere and Endophytic Associations

The rhizosphere and endosphere are major ecological niches where intense plant–microbe interactions occur. The rhizosphere refers to the narrow soil zone surrounding plant roots that is strongly influenced by root exudates, microbial activity, and nutrient exchange processes. Root exudates consisting of sugars, amino acids, organic acids, and secondary metabolites act as signaling molecules and nutrient sources that shape rhizosphere microbial communities (Berendsen *et al.*, 2012). This highly dynamic environment supports extensive microbial competition, cooperation, and communication, making the rhizosphere one of the most biologically active regions within soil ecosystems. Rhizosphere microorganisms contribute significantly to nutrient cycling, disease suppression, soil fertility improvement, and stress tolerance in plants. Endophytic microorganisms inhabit internal plant tissues such as roots, stems, leaves, and seeds without causing disease symptoms. These microorganisms establish either beneficial or neutral relationships with host plants and influence plant metabolism, immunity, and adaptation to environmental stress (Compant *et al.*, 2022). Endophytic bacteria and fungi enhance plant growth through nutrient mobilization, phytohormone production, and protection against pathogens and abiotic stress. The growing understanding of rhizosphere and endophytic associations is contributing significantly to the development of innovative microbial technologies for sustainable and climate-resilient agriculture.

**Table 1. Diversity of plant–microbe interactions in agricultural system**


Interaction Type	Microbial Group	Plant Response	Examples
Symbiosis	Rhizobium	Nitrogen fixation	Legume-rhizobia symbiosis
	AMF	Nutrient uptake, Stress tolerance	Mycorrhizal symbiosis
	Frankia	Nitrogen fixation	Actinorrhizal symbiosis
Commensalism	Plant growth-promoting fungi	Altered root exudation	Bacterial colonization of the rhizosphere
	Nitrogen-fixing cyanobacteria	Nitrogen fixation, growth promotion	Cyanobacterial-plant symbiosis
Mutualism	Plant growth-promoting rhizobacteria (PGPR)	Growth promotion, stress resistance	Rhizobacteria-plant symbiosis
	Mycorrhizal fungi	Nutrient acquisition, disease resistance	Mycorrhizal symbiosis
	Endophytic bacteria	Disease resistance, growth promotion	Endophytic bacteria-plant symbiosis
Amensalism	Allelopathic microorganisms	Inhibition of competing plants	Allelopathy in soil microbial communities
Competition	Soil bacteria and fungi	Nutrient competition, antibiotic production	Fungal colonization of the rhizosphere
	Root pathogens	Disease development, reduced growth	Pathogen-plant interactions in the rhizosphere
Antibiosis	Antibiotic-producing bacteria	Suppression of pathogens, pests	Antibiosis in soil microbiome
	Antifungal-producing fungi	Suppression of fungal pathogens	Fungal-plant interactions
	Allelopathic plants	Inhibition of microbial growth	Plant–microbe interactions in allelopathic systems

Adapted from [Tharanath et al. \(2024\)](#)

### Beneficial Microorganisms and their Functions

Beneficial microorganisms are essential components of sustainable agricultural systems because of their significant contributions to plant nutrition, soil fertility, stress tolerance, and disease suppression. These microorganisms establish mutualistic or associative relationships with plants and influence multiple biological processes within the rhizosphere and plant tissues. In modern agriculture, increasing concerns

regarding excessive use of chemical fertilizers and pesticides have accelerated interest in microbial-based alternatives capable of improving crop productivity while minimizing environmental degradation. Beneficial microorganisms such as nitrogen-fixing bacteria, mycorrhizal fungi, plant growth-promoting rhizobacteria, and microbial biocontrol agents are now recognized as critical tools for environmentally sustainable and climate-resilient agriculture ([Berendsen et al., 2012](#)).



**Figure 2.** Schematic illustration of soil microbes, their roles and applications in agriculture and environment. The figure illustrates the roles of soil microorganisms in nutrient cycling, plant growth promotion, disease suppression, and soil health improvement, as well as their applications in sustainable agriculture and environmental management.

Source: Kiprotich *et al.* (2025)

### Nitrogen-Fixing Bacteria

Nitrogen is a fundamental nutrient required for plant growth and metabolic activities, yet atmospheric nitrogen cannot be directly utilized by most plants. Nitrogen-fixing bacteria possess specialized enzymatic systems capable of converting atmospheric nitrogen into ammonia through biological nitrogen fixation. Symbiotic bacteria such as *Rhizobium* establish highly specialized relationships with leguminous plants by forming root nodules where nitrogen fixation occurs (Lindström and Mousavi, 2019). This symbiotic interaction significantly improves nitrogen availability in soils while reducing dependence on synthetic nitrogen fertilizers. In addition to symbiotic forms, free-living and associative nitrogen-fixing bacteria including *Azospirillum* and *Azotobacter* contribute substantially to soil fertility and crop productivity. These microorganisms improve nutrient availability, stimulate root growth, and enhance plant physiological performance under environmental stress conditions (Mahmud *et al.*, 2020). Nitrogen-fixing microorganisms also contribute to sustainable agricultural productivity by improving soil biological activity and maintaining long-term ecosystem stability. Their increasing application as biofertilizers reflects growing efforts to develop environmentally friendly nutrient management strategies within modern agricultural systems.

### Mycorrhizal Fungi

Mycorrhizal fungi are among the most ecologically important beneficial microorganisms associated with plants. These fungi

establish mutualistic relationships with plant roots through specialized fungal structures that facilitate nutrient and water exchange between plants and fungal hyphae. Arbuscular mycorrhizal fungi (AMF) are particularly significant because they colonize the roots of most terrestrial plant species and improve nutrient acquisition, especially phosphorus uptake in nutrient-deficient soils (Chen *et al.*, 2018). Through extensive hyphal networks, mycorrhizal fungi increase root absorptive capacity and improve plant access to water and essential mineral nutrients. Beyond nutrient acquisition, mycorrhizal fungi contribute significantly to plant adaptation under abiotic stress conditions such as drought, salinity, and heavy metal toxicity. These fungi regulate plant physiological and biochemical processes that improve osmotic balance, antioxidant activity, and stress-responsive signaling pathways (Begum *et al.*, 2019). Mycorrhizal associations also improve soil aggregation and enhance soil structural stability, thereby contributing to long-term soil health and ecosystem sustainability. Their multifunctional roles make mycorrhizal fungi highly valuable for sustainable crop production and ecological restoration programs.

### Plant Growth-Promoting Microorganisms

Plant growth-promoting microorganisms (PGPM) comprise diverse groups of bacteria and fungi capable of enhancing plant growth through direct and indirect mechanisms. Among these, plant growth-promoting rhizobacteria (PGPR) are extensively studied because of their strong influence on root development, nutrient mobilization, and plant physiological regulation (Vejan *et al.*, 2016). Important PGPR genera

include *Bacillus*, *Pseudomonas*, *Azospirillum*, and *Enterobacter*, many of which produce phytohormones such as auxins, cytokinins, and gibberellins that stimulate plant growth and root architecture. Plant growth-promoting rhizobacteria also improve plant tolerance to environmental stresses through regulation of antioxidant systems, osmotic adjustment, and stress-responsive metabolites. Under drought conditions, PGPR enhance water uptake efficiency and reduce oxidative stress, thereby improving crop resilience and productivity (Ojuederie *et al.*, 2019). Furthermore, microbial consortia involving multiple beneficial microorganisms often produce synergistic effects that enhance nutrient use efficiency and plant performance beyond the effects of individual strains (Olanrewaju and Babalola, 2019). The increasing understanding of underground microbial interactions has further emphasized the importance of PGPR in maintaining plant health and rhizosphere stability within sustainable agricultural ecosystems (Bhat *et al.*, 2023).

### Biocontrol Agents

Biocontrol microorganisms are beneficial microbes capable of suppressing plant pathogens and reducing disease incidence through natural biological mechanisms. These microorganisms

include bacteria, fungi, and actinomycetes that inhibit pathogens through competition for nutrients and ecological niches, secretion of antimicrobial compounds, production of lytic enzymes, and induction of plant systemic resistance (Köhl *et al.*, 2019). Biocontrol agents therefore provide environmentally sustainable alternatives to chemical pesticides and contribute significantly to integrated disease management strategies. Fungal species such as *Trichoderma* are widely recognized for their strong biocontrol capabilities and additional plant growth-promoting effects. Certain endophytic strains of *Trichoderma* improve photosynthetic efficiency, nutrient acquisition, and stress tolerance while simultaneously suppressing plant pathogens (Harman *et al.*, 2021). Beneficial rhizosphere microorganisms also strengthen plant immune responses and reduce pathogen colonization within plant tissues. In contrast, pathogenic microorganisms such as *Pseudomonas syringae* possess specialized virulence mechanisms that enable successful host infection and immune suppression (Xin *et al.*, 2018). Understanding these contrasting microbial interactions is essential for developing effective biological control technologies and sustainable crop protection systems.

**Table 2. Beneficial soil microorganisms and their functions**

Types of microorganism	Example species	Benefits
Bacteria	<i>Rhizobium</i> spp.	Nitrogen fixation, enhances plant growth
Fungi	Mycorrhizal Fungi	Improve nutrient and water uptake, promote plant growth
Protozoa	Amoebae, flagellates, ciliate	Regulate bacterial population, nutrients cycling
Algae	Cyanobacteria (blue green algae)	Nitrogen fixation, soil stabilization

Adapted from Pandey (2023)

### Mechanisms and Agricultural Applications of Plant–Microbe Interactions

Plant–microbe interactions are fundamental biological processes that regulate nutrient cycling, plant development, stress adaptation, and soil ecosystem stability within agricultural systems. Beneficial microorganisms inhabiting the rhizosphere and plant tissues establish highly dynamic associations with plants through biochemical signaling, metabolic exchange, and ecological interactions. These mechanisms have gained substantial agricultural importance because they provide sustainable alternatives to excessive dependence on synthetic fertilizers and pesticides, whose prolonged use negatively affects soil biodiversity, beneficial microorganisms, and environmental health (Pandey, 2023). Consequently, microbial-mediated processes are increasingly recognized as essential components of sustainable and climate-resilient agriculture (Compant *et al.*, 2025).

One of the most important mechanisms of plant–microbe interactions is nutrient mobilization within the rhizosphere.

Plant growth-promoting rhizobacteria (PGPR) and other beneficial microorganisms enhance nutrient availability through biological nitrogen fixation, phosphate solubilization, potassium mobilization, and siderophore-mediated iron acquisition. These microbial activities improve nutrient uptake efficiency and support plant growth under nutrient-limited conditions (Vejan *et al.*, 2016). Nitrogen-fixing bacteria are particularly important because they convert atmospheric nitrogen into plant-accessible forms, thereby reducing dependence on synthetic nitrogen fertilizers and improving soil fertility (Liu-Xu *et al.*, 2024). Root exudates further regulate microbial recruitment and activity around plant roots, enabling plants to establish selective associations with beneficial microbial communities capable of improving nutrient acquisition and rhizosphere functionality (Alzate Zuluaga *et al.*, 2024). These interactions form the biological foundation of microbial biofertilizers increasingly used in sustainable crop production systems.


Beneficial microorganisms also contribute substantially to plant growth promotion and tolerance to environmental stress.

Rhizospheric bacteria, fungal endophytes, and mycorrhizal fungi produce phytohormones such as auxins, gibberellins, cytokinins, and ACC deaminase enzymes that regulate root architecture, seed germination, nutrient absorption, and overall plant physiological performance (Nadarajah and Abdul Rahman, 2023). Under adverse environmental conditions including drought, salinity, and temperature stress, beneficial microorganisms enhance osmotic regulation, antioxidant activity, and stress-responsive signaling pathways that improve plant adaptation and survival. Endophytic microorganisms further improve crop resilience by stabilizing plant metabolic processes and enhancing water and nutrient uptake efficiency during environmental stress conditions (Sena *et al.*, 2024). These microbial-mediated mechanisms are increasingly important for maintaining agricultural productivity under changing climatic conditions.

Another major mechanism involves microbial suppression of plant pathogens and regulation of plant immune responses. Beneficial microorganisms inhibit pathogens through ecological competition, production of antimicrobial compounds, secretion of hydrolytic enzymes, and activation of induced systemic resistance within host plants. Rhizosphere microbial communities also participate in complex signaling interactions involving root exudates, quorum sensing molecules, and secondary metabolites that collectively regulate microbial colonization and plant defense responses

(Shi *et al.*, 2024). These interactions contribute significantly to the development of microbial biocontrol agents and biopesticides that provide environmentally sustainable alternatives to synthetic pesticides. The increasing application of microbial biocontrol technologies reflects growing efforts to reduce chemical pesticide use while maintaining effective crop protection and ecological stability.

The cumulative effects of nutrient mobilization, growth promotion, stress tolerance, and disease suppression contribute significantly to soil health improvement and sustainable crop productivity. Beneficial microorganisms regulate nutrient cycling, organic matter decomposition, microbial diversity maintenance, and soil aggregation processes that collectively improve soil structure and ecosystem functioning. Mycorrhizal fungal networks and rhizospheric microbial communities enhance soil porosity, water retention capacity, and long-term soil fertility, thereby supporting resilient agricultural systems (Tharanath *et al.*, 2024). Recent advances in plant microbiome research further demonstrate that targeted management of soil microbial communities can improve crop productivity, environmental sustainability, and regenerative agricultural practices (Kiprotich *et al.*, 2025). Harnessing these microbial mechanisms therefore represents a promising strategy for developing biologically integrated agricultural systems capable of supporting future food security and sustainable environmental management.



**Figure 3. Mechanisms of plant–microbe interactions and their roles in sustainable agriculture.** The figure illustrates the major mechanisms governing plant–microbe interactions within the rhizosphere and phyllosphere, highlighting both beneficial and detrimental relationships. It demonstrates how root exudates, microbial signaling, nutrient cycling, and environmental factors influence microbial community dynamics, plant growth, disease resistance, and crop resilience. The diagram further emphasizes the importance of microbial balance in supporting sustainable agricultural productivity and ecosystem stability. Source: [Anas \*et al.\* \(2024\)](#)


## Emerging Research Approaches and Future Directions

Recent advances in plant–microbe interaction research are increasingly driven by integrative technologies that combine molecular biology, computational science, and sustainable agricultural innovation. Modern approaches now extend beyond conventional microbiology toward systems-level investigations of plant microbiomes, enabling deeper understanding of microbial diversity, functionality, and ecological interactions within agricultural ecosystems. These developments are reshaping sustainable agriculture by facilitating targeted microbial management strategies aimed at improving crop productivity, environmental sustainability, and climate resilience (Trivedi *et al.*, 2020). Genomics and metagenomics have revolutionized the study of plant-associated microbial communities by enabling culture-independent characterization of microbial diversity and functional potential. High-throughput sequencing technologies now allow researchers to identify microbial taxa, metabolic pathways, and host–microbe interactions within complex rhizosphere and endophytic environments. Metagenomic analyses have further revealed that microbial community assembly is strongly influenced by plant genotype, soil conditions, and environmental stress factors, thereby providing insights into microbiome dynamics and ecosystem functionality (Müller *et al.*, 2016). Comparative genomic studies have also identified genes associated with microbial colonization, nutrient acquisition, stress adaptation, and pathogen suppression, contributing to the discovery of beneficial microorganisms with agricultural importance (Levy *et al.*, 2018).

Emerging approaches in microbiome engineering and synthetic biology are creating new opportunities for designing microbial consortia capable of improving plant performance under diverse environmental conditions. Synthetic microbial communities are increasingly used to investigate microbial interactions, ecological stability, and functional compatibility within plant-associated microbiomes (Vorholt *et al.*, 2017). In addition, advances in synthetic biology are enabling the modification of microbial metabolic pathways for enhanced nutrient mobilization, stress tolerance, and disease suppression. These approaches support the development of next-generation bioinoculants with greater ecological

adaptability and field performance compared to conventional single-strain microbial products (Compant *et al.*, 2021). Artificial intelligence (AI), machine learning, and precision agriculture technologies are further transforming plant–microbe interaction research by enabling large-scale analysis of biological and environmental datasets. AI-assisted models can predict microbial behavior, soil health dynamics, crop responses, and disease outbreaks based on genomic, climatic, and agronomic information. Precision agriculture tools including remote sensing, biosensors, and data-driven microbial monitoring systems are increasingly integrated into sustainable farming practices to optimize microbial applications and resource management (Berg *et al.*, 2020). These technologies improve decision-making efficiency and facilitate site-specific agricultural interventions capable of enhancing productivity while minimizing environmental impacts.

Future research is expected to focus strongly on the development of climate-resilient and regenerative agricultural systems supported by beneficial microbiomes. Increasing attention is being directed toward microbial strategies capable of improving crop adaptation to drought, salinity, temperature extremes, and degraded soils under climate change scenarios. Regenerative agricultural practices that promote soil biodiversity, carbon sequestration, and ecological restoration are also expected to accelerate the application of microbiome-based technologies in sustainable food production systems (Philippot *et al.*, 2013). Furthermore, advances in multi-omics integration, microbiome engineering, and predictive computational modeling are likely to support the emergence of highly specialized microbial products tailored for specific crops, environments, and stress conditions. Collectively, these emerging research approaches highlight the growing importance of plant microbiome science in shaping the future of sustainable and resilient agriculture.



**Figure 4. Future directions in plant-microbe interactions include multi-omics, microbial consortia, field applications, climate impacts, emerging pathogens, and sustainable practices. This figure outlines a strategic framework to advance plant-microbe interactions under climate stress, focusing on sustainable agricultural practices**

Source: [Anas et al. \(2024\)](#)

## Conclusion

Plant-microbe interactions play a fundamental role in maintaining plant health, soil fertility, and ecosystem sustainability within agricultural systems. The growing understanding of rhizosphere and endophytic microbial communities has revealed their importance in nutrient mobilization, growth promotion, stress tolerance, and disease suppression. These interactions provide environmentally sustainable alternatives to excessive agrochemical dependence and contribute significantly to improving crop productivity and resilience under changing environmental conditions. Recent advances in genomics, metagenomics, microbiome engineering, synthetic biology, and precision agriculture have transformed plant-microbe research from descriptive studies into highly integrated and application-oriented scientific fields. The development of microbial biofertilizers, biopesticides, and climate-resilient microbial technologies demonstrates the increasing potential of beneficial microorganisms in sustainable agricultural production systems. Furthermore, the integration of microbiome-based approaches into regenerative and climate-smart agriculture offers promising opportunities for restoring soil health, enhancing biodiversity, and supporting long-term food security.

Despite these advances, several challenges remain regarding field-level consistency, ecological adaptability, commercialization, and regulatory standardization of microbial products. Addressing these limitations will require interdisciplinary collaboration among microbiologists, agronomists, biotechnologists, environmental scientists, and policymakers. Future research should focus on understanding complex microbial interactions under diverse environmental conditions and developing highly efficient microbial formulations tailored to specific crops and ecosystems.

Overall, harnessing beneficial plant-microbe interactions represents a transformative strategy for achieving sustainable, resilient, and environmentally responsible agriculture. Continued scientific innovation and practical implementation of microbiome-based technologies will play an increasingly important role in supporting global agricultural sustainability and food production in the face of future climatic and ecological challenges.

## Author Contributions

The author has significantly contributed to the study's conception, design, data collection, analysis, and interpretation. All authors were involved in writing the manuscript or critically reviewing it for its intellectual value. They have reviewed and approved the final version for submission and publication and accept full responsibility for the content and integrity of the work.

## Financial Support

The authors declare that no external funding was received for this study.

## Conflicts of Interest

The authors state that there are no financial, personal, or professional conflicts of interest associated with this research.

## Ethical Approvals

This research did not involve the use of human participants or animal subjects and therefore did not require ethical clearance

## Publisher's Note

The Journal stays neutral about jurisdictional claims in published maps and institutional affiliations

## References

- Alzate Zuluaga MY, Fattorini R, Cesco S and Pii Y (2024) Plant-microbe interactions in the rhizosphere for smarter and more sustainable crop fertilization: the case of PGPR-based biofertilizers. *Front. Microbiol.* 15:1440978. [10.3389/fmicb.2024.1440978](https://doi.org/10.3389/fmicb.2024.1440978)
- Anas, M., Khalid, A., Saleem, H., Khan, K., Khattak, A. and Sh, F. (2024). Symbiotic Synergy: Unveiling Plant-Microbe Interactions in Stress Adaptation. *Journal of Crop Health.* 77(1):18. DOI:10.1007/s10343-024-01070-z
- Backer, R., Rokem, J.S., Ilangumaran, G., Lamont, J., Praslickova, D., Ricci, E. *et al.* (2018) Plant Growth-Promoting Rhizobacteria: Context, Mechanisms of Action, and Roadmap to Commercialization of Biostimulants for Sustainable Agriculture. *Front. Plant Sci.* 9:1473. doi: 10.3389/fpls.2018.01473
- Bais, H., Prithiviraj, B., Jha, A. *et al.* (2005). Mediation of pathogen resistance by exudation of antimicrobials from roots. *Nature* 434, 217–221. <https://doi.org/10.1038/nature03356>
- Begum, N., Qin, C., Ahanger, M., Raza, S., Khan, M., Ashraf, M. *et al.* (2019) Role of Arbuscular Mycorrhizal Fungi in Plant Growth Regulation: Implications in Abiotic Stress Tolerance. *Front. Plant Sci.* 10:1068. doi: 10.3389/fpls.2019.01068
- Ben, J.J. Lugtenberg, R., Caradus, L. J. (2016). Fungal endophytes for sustainable crop production, *FEMS Microbiology Ecology.* 92(12): fiw194, <https://doi.org/10.1093/femsec/fiw194>
- Berendsen, R. L., Pieterse, C. M. and Bakker, P. A. (2012). The rhizosphere microbiome and plant health. *Trends in plant science.* 17(8): 478–486. <https://doi.org/10.1016/j.tplants.2012.04.001>
- Berg, G., Rybakova, D., Fischer, D. *et al.* (2020). Microbiome definition re-visited: old concepts and new challenges. *Microbiome.* 8: 103. <https://doi.org/10.1186/s40168-020-00875-0>
- Berg, G., Rybakova, D., Fischer, D. *et al.* (2020). Microbiome definition re-visited: old concepts and new challenges. *Microbiome.* 8: 103. <https://doi.org/10.1186/s40168-020-00875-0>
- Bhat, M., Mishra, A.K., Jan, S., Bhat, M., Kamal, M., Rahman, S. *et al.* (2023). Plant Growth Promoting Rhizobacteria in Plant Health: A Perspective Study of the Underground Interaction. *Plants (Basel).* 31;12(3):629. doi: 10.3390/plants12030629. PMID: 36771713; PMCID: PMC9919780.
- Chandra, P., Dhuli, P., Verma, P., Singh, A. *et al.* (2020). Culturable microbial diversity in the rhizosphere of different biotypes under variable salinity. *Tropical Ecology.* 61(9):3. DOI:10.1007/s42965-020-00089-3
- Chen, M., Arato, M., Borghi, L., Nouri, E. and Reinhardt, D. (2018) Beneficial Services of Arbuscular Mycorrhizal Fungi – From Ecology to Application. *Front. Plant Sci.* 9:1270. doi: 10.3389/fpls.2018.01270
- Compant, S., Cambon, M. C., Vacher, C., Mitter, B., Samad, A. and Sessitsch, A. (2022). The plant endosphere world—Bacterial life within plants. *Environmental Microbiology.* 23(4):1812–1829. <https://doi.org/10.1111/1462-2920.15240>
- Compant, S., Cassan, F., Kostić, T. *et al.* Harnessing the plant microbiome for sustainable crop production. *Nat Rev Microbiol.* 23: 9–23 (2025). <https://doi.org/10.1038/s41579-024-01079-1>
- Compant, S., Samad, A., Faist, H. and Sessitsch, A. (2021). A review on the plant microbiome: Ecology, functions, and emerging trends in microbial application. *Journal of Advanced Research.* 19: 29–37. <https://doi.org/10.1016/j.jare.2019.03.004>
- Compant, S., Samad, A., Faist, H. and Sessitsch, A. (2021). A review on the plant microbiome: Ecology, functions, and emerging trends in microbial application. *Journal of Advanced Research.* 19: 29–37. <https://doi.org/10.1016/j.jare.2019.03.004>
- Finkel, O. M., Castrillo, G., Herrera Paredes, S., Salas González, I. and Dangl, J. L. (2017). Understanding and exploiting plant beneficial microbes. *Current Opinion in Plant Biology.* 38: 155–163. <https://doi.org/10.1016/j.pbi.2017.04.018>
- Genre, A., Lanfranco, L., Perotto, S. *et al.* (2020). Unique and common traits in mycorrhizal symbioses. *Nat Rev Microbiol* 18, 649–660. <https://doi.org/10.1038/s41579-020-0402-3>
- Harman, G.E., Doni, F., Khadka, R.B. and Uphoff, N. (2021). Endophytic strains of *Trichoderma* increase plants' photosynthetic capability, *Journal of Applied Microbiology.* 130(2):529–546, <https://doi.org/10.1111/jam.14368>
- Kiptotich, K., Muema, E., Wekesa, C., Ndombi, S. *et al.* (2025). Unveiling the roles, mechanisms and prospects of soil microbial communities in sustainable agriculture. *Springer Nature. Discover Soil.* 2(1). DOI:10.1007/s44378-025-00037-4
- Köhl, J., Kolnaar, R. and Ravensberg, W.J. (2019) Mode of Action of Microbial Biological Control Agents Against Plant Diseases: Relevance Beyond Efficacy. *Front. Plant Sci.* 10:845. doi: 10.3389/fpls.2019.00845
- Levy, A., Salas Gonzalez, I., Mittelviefhaus, M. *et al.* (2018). Genomic features of bacterial adaptation to plants. *Nat Genet.* 50:138–150. <https://doi.org/10.1038/s41588-017-0012-9>
- Lindström, K. and Mousavi, S. A. (2019). Effectiveness of nitrogen fixation in rhizobia. *Microbial Biotechnology.* 13(5):1314–1335. <https://doi.org/10.1111/1751-7915.13517>
- Liu-Xu, L., González-Hernández, A. I., Camañes, G., Vicedo, B., Scalschi, L. and Llorens, E. (2024). Harnessing Green Helpers: Nitrogen-Fixing Bacteria and Other Beneficial Microorganisms in Plant–Microbe Interactions for Sustainable Agriculture. *Horticulturae.* 10(6):621. <https://doi.org/10.3390/horticulturae10060621>
- Mahmud, K., Makaju, S., Ibrahim, R. and Missaoui, A. (2020). Current Progress in Nitrogen Fixing Plants and Microbiome Research. *Plants.* 9(1):97. <https://doi.org/10.3390/plants9010097>
- Müller, D. B., Vogel, C., Bai, Y. and Vorholt, J. A. (2016). The Plant Microbiota: Systems-Level Insights and Perspectives. *Annual review of genetics.* 50: 211–234. <https://doi.org/10.1146/annurev-genet-120215-034952>
- Nadarajah, K. and Abdul Rahman, N. S. N. (2023). The Microbial Connection to Sustainable Agriculture. *Plants.* 12(12):2307. <https://doi.org/10.3390/plants12122307>
- Ngou, B.P.M., Ahn, H.K., Ding, P. *et al.* (2021). Mutual potentiation of plant immunity by cell-surface and intracellular receptors. *Nature.* 592: 110–115. <https://doi.org/10.1038/s41586-021-03315-7>
- Niu, B., Paulson, J.N., Zheng, X. and Kolter, R. (2017). Simplified and representative bacterial community of maize roots, *Proc. Natl. Acad. Sci. U.S.A.* 114 (12):E2450–E2459. <https://doi.org/10.1073/pnas.1616148114> (2017).
- Ojuederie, O. B., Olanrewaju, O. S. and Babalola, O. O. (2019). Plant Growth Promoting Rhizobacterial Mitigation of Drought Stress in Crop Plants: Implications for Sustainable Agriculture. *Agronomy.* 9(11): 712. <https://doi.org/10.3390/agronomy9110712>
- Olanrewaju, O. and Babalola, O.O. (2019). Bacterial Consortium for Improved Maize (*Zea mays* L.) Production. *Microorganisms.* 7(11):519. doi: 10.3390/microorganisms7110519. PMID: 31683950; PMCID: PMC6920993.

- Oldroyd, G. E. and Dixon, R. (2014). Biotechnological solutions to the nitrogen problem. *Current opinion in biotechnology*. 26:19–24. <https://doi.org/10.1016/j.copbio.2013.08.006>
- Pandey, S. (2023). Impact of Agrochemicals on Beneficial Microorganisms and Human Health. *International Journal of Environment and Climate Change*. 13(10):1135-1145. DOI:10.5281/zenodo.8318889
- Philippot, L., Raaijmakers, J. M., Lemanceau, P. and van der Putten, W. H. (2013). Going back to the roots: the microbial ecology of the rhizosphere. *Nature reviews. Microbiology*. 11(11):789–799. <https://doi.org/10.1038/nrmicro3109>
- Santoyo, G., Guzmán-Guzmán, P., Parra-Cota, F. I., Santos-Villalobos, S. d. I., Orozco-Mosqueda, M. d. C. and Glick, B. R. (2021). Plant Growth Stimulation by Microbial Consortia. *Agronomy*. 11(2):219. <https://doi.org/10.3390/agronomy11020219>
- Sena L, Mica E, Valè G, Vaccino P and Pecchioni N (2024) Exploring the potential of endophyte-plant interactions for improving crop sustainable yields in a changing climate. *Front. Plant Sci*. 15:1349401. doi: 10.3389/fpls.2024.1349401
- Shelake, M., Waghunde, R. and Kim, J. (2026). Coevolution of plant–microbe interactions, friend–foe continuum, and microbiome engineering for a sustainable future. *Molecular Plant*. 19(4): 719-754. <https://doi.org/10.1016/j.molp.2026.01.010>.
- Shi, X., Zhao, Y., Xu, M., Ma, L., Adams, J., Shi, Y. (2024). Insights into plant–microbe interactions in the rhizosphere to promote sustainable agriculture in the new crops era. *New Crops*. 1:100004. <https://doi.org/10.1016/j.ncrops.2023.11.002>
- Tetlow, I. J. and Emes, M. J. (2017). Starch Biosynthesis in the Developing Endosperms of Grasses and Cereals. *Agronomy*. 7(4): 81. <https://doi.org/10.3390/agronomy7040081>
- Tharanath, A. C., Upendra, R. S. and Rajendra, K. (2024). Soil Symphony: A Comprehensive Overview of Plant–Microbe Interactions in Agricultural Systems. *Applied Microbiology*. 4(4): 1549-1567. <https://doi.org/10.3390/applmicrobiol4040106>
- Trivedi, P., Leach, J. E., Tringe, S. G., Sa, T. and Singh, B. K. (2020). Plant-microbiome interactions: from community assembly to plant health. *Nature reviews. Microbiology*. 18(11): 607–621. <https://doi.org/10.1038/s41579-020-0412-1>
- Vejan, P., Abdullah, R., Khadiran, T., Ismail, S. and Nasrulhaq, B. A. (2016). Role of Plant Growth Promoting Rhizobacteria in Agricultural Sustainability-A Review. *Molecules*. 21(5):573. doi: 10.3390/molecules21050573. PMID: 27136521; PMCID: PMC6273255.
- Vorholt, J. A., Vogel, C., Carlström, C. I. and Müller, D. B. (2017). Establishing Causality: Opportunities of Synthetic Communities for Plant Microbiome Research. *Cell host & microbe*. 22(2):142–155. <https://doi.org/10.1016/j.chom.2017.07.004>
- Wang, T., Yang, K., Ma, Q., Jiang, X., Zhou, Y., Kong, D. *et al.* (2022). Rhizosphere Microbial Community Diversity and Function Analysis of Cut Chrysanthemum During Continuous Monocropping. *Frontiers in microbiology*. 13:801546. <https://doi.org/10.3389/fmicb.2022.801546>
- Xin, X. F., Kvitko, B. and He, S. Y. (2018). *Pseudomonas syringae*: what it takes to be a pathogen. *Nature reviews. Microbiology*. 16(5):316–328. <https://doi.org/10.1038/nrmicro.2018.17>
- Zhu, Y., Zhang, Y., Li, X. and colleagues. (2024). Insights into plant–microbe interactions in the rhizosphere to promote sustainable agriculture in the new crops era. *New Crops*. 1: 100004. <https://doi.org/10.1016/j.ncrops.2023.11.002>

#### How to cite this article

Odoh, E. H., Nonye, E. C., Vitus, J. E., Idris, H. S., Abah, M. A., Abimbola, M. O., & Agida, O. D. (2026). Plant-microbe interactions: A review of current research and future directions in sustainable agriculture. *Microbial Science Archives*, 6(2), 25–37. <https://doi.org/10.47587/MSA.2026.6202>

This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

